

## PATIENT INFORMATION

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_ Sex: M / F Ht: \_\_\_\_\_ Wt: \_\_\_\_\_ lbs / kg  
 Primary Language: \_\_\_\_\_ Allergies: \_\_\_\_\_  
 Patient Preferred Location: \_\_\_\_\_

<ICD 10 CODE REQUIRED>

## DIAGNOSIS & CLINICAL INFORMATION

### ICD 10 Code (PROVIDE COMPLETE CODE)

- M08.2 \_\_\_\_\_ Juvenile Rheumatoid Arthritis w/ Systemic Onset
- M08.3 Juvenile Rheumatoid Polyarthritits (Seronegative)
- Other: \_\_\_\_\_

**REQUIRED:** Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.

**LAB RESULTS:** Hepatitis B within 3 years, Negative TB within 12 months. CBC with diff, Platelets, AST, ALT, and Lipid panel within 60 days.

## PRESCRIPTION

### Lab Orders+

#### Required:

Negative TB  
 CBC with diff, Platelets, AST, and ALT, at 2nd infusion, then every 8 weeks for Polyarticular JIA and every 4 weeks for Systemic JIA.  
 Lipid Panel, at 2nd infusion, then every six months  
 +Medix Infusion will draw maintenance labs unless otherwise directed by Referring Provider.

### Actemra (tocilizumab)

Weight < 30 kg infuse in 50 mL of 0.9% Sodium Chloride, weight ≥ 30 kg infuse in 100 mL of 0.9% Sodium Chloride.

- Cytokine Release Syndrome:** every 2 weeks (no < 14 days) for one year  
 IV: (wt < 30 kg): Infuse 12 mg/kg  
 IV: (wt ≥ 30 kg): Infuse 8 mg/kg
- Polyarticular Juvenile Idiopathic Arthritis:** every 4 weeks (no < 28 days) for one year  
 IV: (wt < 30 kg): Infuse 10 mg/kg  
 IV: (wt ≥ 30 kg): Infuse 8 mg/kg
- Systemic Juvenile Idiopathic Arthritis:** every 2 weeks (no < 14 days) for one year  
 IV: (wt < 30 kg): Infuse 12 mg/kg  
 IV: (wt ≥ 30 kg): Infuse 8 mg/kg

Patient Weight: \_\_\_\_\_ lbs or \_\_\_\_\_ kg

In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reaction protocol.

**Post Treatment Observations:** The patient is observed for 30 minutes following the first and second infusion.

**Comments:** \_\_\_\_\_

## PRESCRIBER INFORMATION

Prescriber Name: \_\_\_\_\_ *Signature:* \_\_\_\_\_

Date: \_\_\_\_\_ NPI #: \_\_\_\_\_ Specialty: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ (If Applicable)

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Contact Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Email: \_\_\_\_\_