

PATIENT INFORMATION

Patient Name: _____ DOB: _____ Phone: _____ Sex: M / F Ht: _____ Wt: _____ lbs / kg
 Primary Language: _____ Allergies: _____
 Patient Preferred Location: _____

<ICD 10 CODE REQUIRED>

DIAGNOSIS & CLINICAL INFORMATION

ICD 10 Code (PROVIDE COMPLETE CODE)

DERMATOLOGY

- L40.5 Psoriatic Arthritis/Arthropathy
- L40. Psoriasis

GASTROENTEROLOGY

- K50.0 Crohn's Disease, Small Intestine
- K50.1 Crohn's Disease, Large Intestine

- K50.8 Crohn's Disease, Small & Large Intestine
- K50.9 Crohn's Disease, Unspecified
- K51.8 Other Ulcerative Colitis, Chronic
- K51.5 Left Sided - Ulcerative Colitis, Chronic
- K51.0 Universal Ulcerative Pancolitis, Chronic
- K51.9 Ulcerative Colitis, Unspecified
- K60.3 Anal Fistula
- K63.2 Fistula of Intestine

RHEUMATOLOGY

- M05. Rheumatoid Arthritis, w/ Rheumatoid Factor
- M06. Rheumatoid Arthritis, w/o Rheumatoid Factor
- L40.5 Psoriatic Arthritis/Arthropathy
- M45. Ankylosing Spondylitis
- D86.0 Sarcoidosis of the Lung

OTHER: _____

REQUIRED: Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.
LAB RESULTS: Include Negative Hepatitis B within 3 years and Negative TB within 12 months.

PRESCRIPTION

Pre-Medications

- Acetaminophen: 650 mg PO
- Cetirizine: 10 mg PO
- Diphenhydramine: 25 mg PO
- Diphenhydramine: 25 mg IVP
- Famotidine: 20 mg PO
- Methylprednisolone: 125 mg SIVP
- Other: _____

Lab Orders+

Required: Negative TB, annually
+Medix Infusion will draw maintenance labs unless otherwise directed by Referring Provider.

Renflexis (infliximab-abda)

Infuse in 250 mL of 0.9% NS over at least 2 hours via pump with 0.2-micron filter. Doses > 1000 mg need total volume of 500 mL.

Loading Dose: (SELECT ONE)

- IV:** Infuse 3 mg/kg at weeks 0, 2, and 6
- IV:** Infuse 5 mg/kg at weeks 0, 2, and 6
- IV:** Infuse _____ mg or _____ mg/kg at weeks 0, 2, and 6

Maintenance: (SELECT ONE)

- IV:** Infuse 3 mg/kg every 8 weeks for one year
- IV:** Infuse 5 mg/kg every 8 weeks for one year
- IV:** Infuse _____ mg or _____ mg/kg every _____ weeks for one year

In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reaction protocol.

Post Treatment Observations: The patient is observed for 30 minutes following the first infusion.

Comments: _____

PRESCRIBER INFORMATION

Prescriber Name: _____ **Signature:** _____

Date: _____ NPI #: _____ Specialty: _____

Supervising Physician: _____ (If Applicable)

Address: _____ City: _____ State: _____ Zip: _____

Contact Name: _____ Phone: _____ Fax: _____ Email: _____