

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_ Sex: M F Ht: \_\_\_\_\_ Wt: \_\_\_\_\_ lbs kg  
 Primary Language: \_\_\_\_\_ Allergies: \_\_\_\_\_  
 Patient Preferred Location: \_\_\_\_\_

**<ICD 10 CODE REQUIRED> DIAGNOSIS & CLINICAL INFORMATION**
**ICD 10 Code (PROVIDE COMPLETE CODE)**

M08.2 \_\_\_\_\_ Juvenile Rheumatoid Arthritis w/ Systemic Onset  
 M08.3 Juvenile Rheumatoid Polyarthritis (Seronegative)  
 Other: \_\_\_\_\_

**REQUIRED: Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.**

**LAB RESULTS: Negative TB within 12 months. CBC with diff, Platelets, AST, ALT, and Lipid panel within 60 days.**

**PRESCRIPTION**
**Lab Orders+**
*Required:*

Negative TB  
 CBC with diff, Platelets, AST, and ALT, at 2nd infusion, then every 8 weeks for Polyarticular JIA and every 4 weeks for Systemic JIA.  
 Lipid Panel, at 2nd infusion, then every six months

**+ Medix Infusion will draw maintenance labs unless otherwise directed by Referring Provider**

**Actemra (tocilizumab)**

Weight < 30 kg infuse in 50 mL of 0.9% Sodium Chloride, weight ≥ 30 kg infuse in 100 mL of 0.9% Sodium Chloride.

**Cytokine Release Syndrome:**

every 2 weeks (no < 14 days) for one year

**IV:** (wt < 30 kg): Infuse 12 mg/kg

**IV:** (wt ≥ 30 kg): Infuse 8 mg/kg

**Polyarticular Juvenile Idiopathic Arthritis:**

every 4 weeks (no < 28 days) for one year

**IV:** (wt < 30 kg): Infuse 10 mg/kg

**IV:** (wt ≥ 30 kg): Infuse 8 mg/kg

**Systemic Juvenile Idiopathic Arthritis:**

every 2 weeks (no < 14 days) for one year

**IV:** (wt < 30 kg): Infuse 12 mg/kg

**IV:** (wt ≥ 30 kg): Infuse 8 mg/kg

**Patient Weight:** \_\_\_\_\_ lbs or \_\_\_\_\_ kg

In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reaction protocol.

**Post Treatment Observations:** The patient is observed for 30 minutes following the first and second infusion.

**Comments:**


---



---



---



---

**PRESCRIBER INFORMATION**

Prescriber Name: \_\_\_\_\_ Signature: \_\_\_\_\_  
 Date: \_\_\_\_\_ NPI #: \_\_\_\_\_ Specialty: \_\_\_\_\_  
 Supervising Physician: \_\_\_\_\_ (If Applicable)  
 Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_  
 Contact Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Email: \_\_\_\_\_