## medix infusion

## **Ruxience Order Form**



(rituximab-pvvr)

PATIENT INFORMATION				
Patient Name:  Primary Language:  Patient Preferred Location:	Illergies:			0
<icd 10="" code="" required=""></icd>	DIAGNOSIS & CI	LINICAL INFORMA	TION	
ICD 10 Code M06.9 Rheumatoid Arthritis M31.30 Granulomatosis w/ Polyangitis (V M31.7 Microscopic Polyangitis Other:	/egener's Granulomate	osis GPA)		
<u>REQUIRED</u> : Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy. <u>LAB RESULTS</u> : Include Negative Hepatitis B within 3 years.				
PRESCRIPTION				
Pre-Medications Acetaminophen: 650 mg PO Methylprednisolone: 125 mg SIVP Ruxience (rituximab-pvvr)	Diphenhydramine: 2 Other:	•		
Infuse in 250-550 mL of 0.9% Sodium Chlor	ide			
Dose: (SELECT ONE)				
IV: Infuse 1000 mg Frequency & Duration: (SELECT ONE) Infuse single dose Infuse every week for 4 weeks total Infuse initial dose at day 1 followed by 2m Other frequency:	d dose on day 15, the	n repeat cycle every	nt:, Weight: Ibs	s <b>or</b> kg
In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reaction protocol.				
Post Treatment Observations: The patient is observed for 60 minutes following the first infusion. Comments:				
PRESCRIBER INFORMATION				
Prescriber Name: Date: NPI #: Supervising Physician: Address:		Specialty:		(If Applicable)
Contact Name:	-			