Skyrizi IV Order Form
(risankizumab-rzaa)

medix infusion

FAX TO: 972.499.9210

PATIENT INFORMATION

Patient Name: _____ DOB: _____ Phone: _____ Sex: ☐ M ☐ F Ht: _____ Wt: _____ ☐ lbs ☐ kg

Primary Language: _____ Allergies: _____

Patient Preferred Location: _____

<ICD 10 CODE REQUIRED> DIAGNOSIS & CLINICAL INFORMATION

ICD 10 Code

☐ K50.0 _____ Crohn's Disease, Small Intestine

☐ K50.1 _____ Crohn's Disease, Large Intestine

☐ K50.8 _____ Crohn's Disease, Small & Large Intestine

☐ K50.9 _____ Crohn's Disease, Unspecified

***REQUIRED*: Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.**
***LAB RESULTS*: Include Negative TB within 12 months.**

PRESCRIPTION

Lab Orders+
Required: Negative TB,
Liver enzymes and bilirubin at weeks 0 and 4
+Medix Infusion will draw maintenance labs unless otherwise directed by Referring Provider.

Skyrizi IV (risankizumab-rzaa)

Loading Dose IV:
Infuse 600 mg in 250 ml of 5% Dextrose over at least 1 hour at weeks 0, 4, and 8

In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reaction protocol.

Post Treatment Observations: The patient is observed for 30 minutes following the first infusion.

Comments:

_____

_____

_____

_____

_____

PRESCRIBER INFORMATION

Prescriber Name: _____ Signature: _____

Date: _____ NPI #: _____ Specialty: _____

Supervising Physician: _____ (If Applicable)

Address: _____ City: _____ State: _____ Zip: _____

Contact Name: _____ Phone: _____ Fax: _____ Email: _____