medix infusion

Krystexxa Order Form (pegloticase)



PATIENT INFORMATION			
Patient Name: Primary Language: Allergi Patient Preferred Location:	DOB: Phone: es:	Sex: M F Ht:Wt:	Ū
<pre></pre> <pre></pre> <pre></pre> <pre>diagnosis & Clinical Information</pre>			
ICD 10 Code (PROVIDE COMPLETE CODE) M1A0 Chronic Gout, w/o Tophi M1A1 Chronic Gout, w/ Tophi Other:	It is recommended that the patient discontinue oral urate-lowering medications 2-3 days (up to one week) before starting Krystexxa.		
<u>REQUIRED</u> : Demographics and Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to in- clude any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy. <u>LAB RESULTS</u> : : G6PD, baseline uric acid > 6.0 mg/dL.			
PRESCRIPTION			
Pre-Medications Required: Acetaminophen: 650 mg PO, may repeat q 4-6 hours, PRN infusion reaction Diphenhydramine: 25 mg IVP, may repeat q 6 hours, PRN infusion reaction Methylprednisolone: 125 mg SIVP Other: KRYSTEXXA (pegloticase)		Lab Orders+ Required: Uric Acid Level, 24-72 hours prior to infusion If Uric Acid Level > 6 mg/dL upon two consecutive lab draws, hold dose, and contact prescriber +Medix Infusion will draw maintenance labs unless otherwise directed by Referring Provider	
Loading Dose			
 IV: Infuse 8 mg in 250 mL of 0.9% Sodium Chloride over 2 hours, every 2 weeks for one year Post Treatment Observations: The patient is observed for 60 minutes following the first and second administrations. 			
Adverse Events: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.			
PRESCRIBER INFORMATION			
Prescriber Name:	Sig	nature:	
Date: NPI #:	-	-	
Supervising Physician:			(If Applicable)
Address:	City:	State: Zip:_	

__ Fax: __

_ Phone: ___

Contact Name: _

Email: _