

Nulojix Order Form (belatacept)

FAX TO: 972.499.9210

PATIENT INFORMATION								
Patient Name:	DOB:	_ Phone:	Sex:	M F	Ht:	_ Wt:	_ lbs	kg
Primary Language:	Allergies:							
Patient Preferred Location:								
<icd 10="" code="" required=""> DIAGNOSIS & CLINICAL INFORMATION</icd>								
ICD 10 Code Z94.0 Kidney Transplant Status Other:		Prescribing Information Nulojix is contraindicated in transplant recipients who are Epstein-Barr (EBV) seronegative or have an unknown serostatus.						S .
		Patient <u>MUST</u> be enrolled in the Nulojix Distribution Program (NDP) and have a patient ID number from NDP. Medication <u>cannot</u> be ordered for new or exisiting patients without ID number. Call Bristol-Myers Squibb at 855.511.6180 to enroll.						
<u>REQUIRED</u> : Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy. <u>LAB RESULTS:</u> Include Negative TB and Epstein-Barr serology. PRESCRIPTION								
Nulojix Distribution Program Patient ID #: _ Date of Patient's Last Dose of Nulojix: Transplant Date:		Lab Orders+ Required: Negative TB, anually						
Weight at Transplant: lbs or _ Patient Current Weight: lbs or * Dose is calculated on transplant weigh	kg r kg		+Medix Infusion supplied by R				BMP if	not
NULOJIX (belatacept)								
Loading Dose								
IV: Infuse 5 mg/kg in 100 mL of 0.19% Sod of 30 minutes via pump using a 0.2-micron			Duration:					
Post Treatment Observations: The patient is observed for 30 minutes following the first administration.								
Adverse Events: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol. Comments:								
PRESCRIBER INFORMATION								
Prescriber Name:		Signature: _						
Date: NPI #:		Specialty:						
Supervising Physician:						(If	Applica	ıble)
Address:	City:			State:		Zip:		_
Contact Name:	Phone:	Fax:		Email: _				