

Ocrevus Order Form (ocrelizumab)

FAX TO: 972.499.9210

PATIENT INFORMATION				
Patient Name:	DOB:	Phone:	Sex: M F Ht:	Wt: lbs kg
Primary Language:	Allergies:			
Patient Preferred Location:				
<icd 10="" code="" required=""> DIAGNOSIS & CLINICAL INFORMATION</icd>				
ICD 10 Code G35 Relapsing Remitting Multiple Scleros G35 Primary Progressive Multiple Scleros Other:				
<u>REQUIRED</u> : Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.				
<u>LAB RESULTS:</u> Include Negative Hepatitis B within 3 years to initiate therapy.				
PRESCRIPTION				
Pre-Medications Required: Acetaminophen: 500 mg PO, may repeat q	4-6 hours, PRN infus	ion reaction		
Select Route: Diphenhydramine: 25 mg PO, may repeat q 6 hours, PRN infusion reaction Diphenhydramine: 25 mg IVP, may repeat q 6 hours, PRN infusion reaction Methylprednisolone: 125 mg SIVP Other:				
OCREVUS (ocrelizumab)				
Loading Dose IV: Infuse 300 mg in 250 mL of 0.9% Sodium Chloride over at least 2 hours and 30 minutes via pump using a 0.2-micron filter at weeks 0 and 2				
Maintenance Dose (FROM WEEK 0) IV: Infuse 600 mg in 500 mL of 0.9% Sodium Chloride over 2 hours or longer via pump using a 0.2 micron filter every 6 months for one year				
Post Treatment Observations: The patient is observed for 60 minutes following the first administration.				
Adverse Events: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.				
Comments:				
PRESCRIBER INFORMATION				
Prescriber Name:		Signature:		
Date: NPI #:				
Supervising Physician:				(If Applicable)
Address:	City:		State:	Zip:
Contact Name:	Phone:	Fax:	Email:	