

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_ Sex: M F Ht: \_\_\_\_\_ Wt: \_\_\_\_\_ lbs kg

Primary Language: \_\_\_\_\_ Allergies: \_\_\_\_\_

**Patient Preferred Location:** \_\_\_\_\_**<ICD 10 CODE REQUIRED> DIAGNOSIS & CLINICAL INFORMATION****ICD 10 Code (PROVIDE COMPLETE CODE)**

ICD 10 Code: \_\_\_\_\_

Description: \_\_\_\_\_

**REQUIRED: Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.****LAB RESULTS: CMP or AST/ALT within last 90 days. Include culture report.****PRESCRIPTION****REZZAYO (rezafungin)****Lab Orders+****Loading Dose**

IV: Infuse 400 mg in 250 mL of 0.9% Sodium Chloride over 60 minutes

**+ Medix Infusion will draw required CBC/AST/ALT if not supplied by Referring Provider****Maintenance Dose**

IV: Infuse 200 mg in 250 mL of 0.9% Sodium Chloride over 60 minutes weekly for \_\_\_\_\_ weeks

Maintenance dosing to start 1 week following initial dose

**Post Treatment Observations:** The patient is observed for 30 minutes following the first and second administrations.**Adverse Events:** In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.**Comments:**\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_**PRESCRIBER INFORMATION**

Prescriber Name: \_\_\_\_\_ Signature: \_\_\_\_\_

Date: \_\_\_\_\_ NPI #: \_\_\_\_\_ Specialty: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ (If Applicable)

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Contact Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Email: \_\_\_\_\_