

Rituximab Form

(rituximab)

FAX TO: 972.499.9210

PATIENT INFORMATION				
Patient Name:	DOB:	_ Phone: Se	x: M F Ht: W	t: lbs kg
Primary Language:	Allergies:			
Patient Preferred Location:				
<icd 10="" code="" required=""> DIAGNOSIS & CLINICAL INFORMATION</icd>				
ICD 10 Code M06.9 Rheumatoid Arthritis M31.30 Granulomatosis w/Polyangitis (WM31.7 Microscopic Polyangitis Other:		,		
<u>REQUIRED</u> : Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy. <u>LAB RESULTS:</u> Include Negative Hepatitis B within 3 years.				
	PRE	SCRIPTION		
Pre-Medications				
Acetaminophen: 650 mg PO Methylprednisolone: 125 mg SIVP				
RITUXIMAB (rituximab)				7
Rituxan (rituximab) or Biosimilar as dictar Medix Infusion will determine apporpr OR Rituximab product Infuse in 250-550 ml of 0.9% Sodium Ch	riate product pbased (DO NO	upon benefit investigation	MEDIX USE ONLY Product to be Used: Rituxan Truxima Ruxience	
Loading Dose (SELECT ONE) IV: Infuse 1000 mg	IV: infuse 375 mg	/m² – Required → Height:	Weight: lbs	or kg
Frequency and Duration (SELECT ONE) Infuse Single Dose Infuse every week for 4 weeks total Infuse initial dose at day 1 followed by 2nd dose on day 15, then repeat cycle every months for one year Other frequency: for one year				
Post Treatment Observation: The patient is observed for 60 minutes following the first administration.				
Adverse Reactions: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.				
Comments:				
PRESCRIBER INFORMATION				
Prescriber Name:		Signature:		
Date: NPI #:		Specialty:		
Supervising Physician:				(If Applicable)
Address:	City:		_ State:	Zip:
Contact Name:	Phone:	Fax:	Email:	