

## **Ruxience Order Form**

(rituximab-pvvr)

FAX TO: 972.499.9210

PATIENT INFORMATION					
Patient Name:	DOB:	_ Phone:	Sex: M	1 F Ht:	_Wt: lbs kg
Primary Language:	Allergies:				
Patient Preferred Location:					
<icd 10="" code="" required=""> DIAGNOSIS &amp; CLINICAL INFORMATION</icd>					
ICD 10 Code  M06.9 Rheumatoid Arthritis  M31.30 Granulomatosis w/Polyangitis (Wegener's Granulomatosis GPA)  M31.7 Microscopic Polyangitis  Other:					
<u>REQUIRED</u> : Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy. <u>LAB RESULTS:</u> Include Negative Hepatitis B within 3 years.					
PRESCRIPTION					
Pre-Medications Acetaminophen: 650 mg PO Methylprednisolone: 125 mg SIVP	Diphenhydramine:				
RUXIENCE (rituximab-pvvr)					
Infuse in 250-550 mL of 0.9% Sodium Chloride					
Loading Dose (SELECT ONE)					
IV: Infuse 1000 mg	IV: Infuse 375 mg/	/m² – <b>Required</b> → Height:	, We	eight:	lbs <b>or</b> kg
Infuse single dose Infuse every week for 4 weeks total Infuse initial dose at day 1 followed by 2nd dose on day 15, then repeat cycle every months for one year Other frequency: for one year					
Post Treatment Observations: The patient is observed for 60 minutes following the first administration.					
Adverse Events: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.					
Comments:					
PRESCRIBER INFORMATION					
Prescriber Name: Signature:					
Date: NPI #:		-			
Supervising Physician:					(If Applicable)
Address:	City:		State	e:	Zip:
Contact Name:	Phone:	Fax:	Em	nail:	