

## PATIENT INFORMATION

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_ Sex: M F Ht: \_\_\_\_\_ Wt: \_\_\_\_\_ lbs kg

Primary Language: \_\_\_\_\_ Allergies: \_\_\_\_\_

Patient Preferred Location: \_\_\_\_\_

## <ICD 10 CODE REQUIRED> DIAGNOSIS & CLINICAL INFORMATION

### ICD 10 Code

D58.8 Other Specified Hereditary Hemolytic Anemias  
D59.3 Hemolytic Uremic Syndrome  
D59.4 Other Non-Autoimmune Hemolytic Anemias  
(Including Microangiopathic Hemolytic Anemia)  
D59.5 Paroxysmal Nocturnal Hemoglobinuria  
D59.8 Other Acquired Hemolytic Anemias

### ICD 10 Code

G36.0 Neuromyelitis Optica  
G70.00 Generalized Myasthenia Gravis, w/o Acute Exacerbation  
G70.01 Generalized Myasthenia Gravis, w/Acute Exacerbation  
Other: \_\_\_\_\_

### Prescribing Information

Meningococcal document required for all diagnoses. See Pre-Medications and Required Labs by diagnosis below.

**REQUIRED:** Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.  
**LAB RESULTS:** Include labs to support diagnosis.

## PRESCRIPTION

### SOLIRIS (eculizumab)

Administer over at least 35 minutes in adults, not to exceed 2 hours.

### PAROXYSMAL NOCTURNAL HEMOGLOBINURIA (PNH)

#### Loading Dose

IV: Infuse 600 mg dose weekly for first 4 weeks followed by 900 mg dose at week 5

#### Maintenance Dose

IV: Infuse 900 mg dose every 2 weeks for one year

### ATYPICAL HEMOLYTIC UREMIC SYNDROME (aHUS)

#### Loading Dose

IV: Infuse 900 mg dose weekly for first 4 weeks followed by 1200 mg dose at week 5

#### Maintenance Dose

IV: Infuse 1200 mg dose every 2 weeks for one year

### GENERALIZED MYASTHENIA GRAVIS (gMG) and NEUROMYELITIS OPTICA SPECTRUM DISORDER (NMOSD)

#### Loading Dose

IV: Infuse 900 mg dose weekly for first 4 weeks followed by 1200 mg dose at week 5

#### Maintenance Dose

IV: Infuse 1200 mg dose every 2 weeks for one year

### Pre-Medications

Acetaminophen: 650 mg PO  
Cetirizine: 10 mg PO  
Diphenhydramine: 25 mg PO  
Diphenhydramine: 25 mg IVP  
Other: \_\_\_\_\_

### Pre-Medications

Acetaminophen: 650 mg PO  
Cetirizine: 10 mg PO  
Diphenhydramine: 25 mg PO  
Diphenhydramine: 25 mg IVP  
Other: \_\_\_\_\_

### Pre-Medications

Acetaminophen: 650 mg PO  
Cetirizine: 10 mg PO  
Diphenhydramine: 25 mg PO  
Diphenhydramine: 25 mg IVP  
Other: \_\_\_\_\_

SOLIRIS	DILUENT VOLUME	FINAL VOLUME
300 mg	30 mL	60 mL
600 mg	60 mL	120 mL
900 mg	90 mL	180 mL
1200 mg	120 mL	240 mL

### Required Labs

- Baseline Serum Lactate Dehydrogenase (LHD) Hemoglobin Level
- Documented Meningococcal Vaccine

### Required Labs

- Baseline Serum Lactate Dehydrogenase (LHD) Hemoglobin Level
- Serum Creatinine/eGFR
- Platelet Count
- Plasma Exchange
- Documented Meningococcal Vaccine

### Required Labs

- Positive Serologic Test for Anti-AChR Antibodies
- Documented Meningococcal Vaccine

**Post Treatment Observations:** The patient is observed for 60 minutes following the first administration.

**Adverse Events:** In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.

## PRESCRIBER INFORMATION

Prescriber Name: \_\_\_\_\_ Signature: \_\_\_\_\_

Date: \_\_\_\_\_ NPI #: \_\_\_\_\_ Specialty: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ (If Applicable)

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Contact Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Email: \_\_\_\_\_