

Spevigo Order Form (spesolimab-sbzo)

FAX TO: 972.499.9210

		PATIEI	NT INFORMATION		
Dationt Non-s				Sex: M F Ht:	\\\/\tag{\tag{\tag{\tag{\tag{\tag{\tag{
					9
		_			
Patient Preferred Location:					
<icd 10="" code="" requi<="" td=""><td>RED></td><td>DIAGNOSIS &</td><td>CLINICAL INFOR</td><td>MATION</td><td></td></icd>	RED>	DIAGNOSIS &	CLINICAL INFOR	MATION	
ICD 10 Code L40.1 Generalized pu	istular psoriasis				
<u>REQUIRED</u> : Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy. <u>LAB RESULTS:</u> Include Negative Hepatitis TB within 12 months.					
PRESCRIPTION					
Dro Modications			Lab Ordara		
Pre-Medications			<u>Lab Orders</u> + Required: Nega	tive TB	
Acetaminophen: 650 Diphenhydramine: 25			,		
Diphenhydramine: 25 mg IVP Other:		 + Medix Infusion will draw maintenance labs unless otherwis by Referring Provider 			ess otherwise directed
Outer:					
SPEVIGO (spesolimab)					
<u>Loading Dose</u> IV: Infuse 900 mg in 100 mL of 0.9% Sodium Chloride over 90 minutes via pump with 0.2-micron filter					
Repeat dose 7 days following initial dose if flare persists					
Post Treatment Observation: The patient is observed for 30 minutes following the first administration.					
Adverse Reactions: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reaction protocol.					
Comments:					
PRESCRIBER INFORMATION					
Prescriber Name:			Signature: _		
Date:	NPI #:		Specialty:		
Supervising Physician:					(If Applicable)
Address:		City:		State:	Zip:
Contact Name:		Phone:	Fax:	Email:	