

PATIENT INFORMATION

Patient Name: _____ DOB: _____ Phone: _____ Sex: M F Ht: _____ Wt: _____ lbs kg

Primary Language: _____ Allergies: _____

Patient Preferred Location: _____

<ICD 10 CODE REQUIRED> DIAGNOSIS & CLINICAL INFORMATION

ICD 10 Code (PROVIDE COMPLETE CODE)

- | | |
|--|---|
| K50.0 _____ Crohn's Disease, Small Intestine | K51.8 _____ Other Ulcerative Colitis, Chronic |
| K50.1 _____ Crohn's Disease, Large Intestine | K51.5. _____ Left Sided - Ulcerative Colitis, Chronic |
| K50.8 _____ Crohn's Disease, Small & Large Intestine | K51.0. _____ Universal Ulcerative Pancolitis, Chronic |
| K50.9 _____ Crohn's Disease, Unspecified | K51.9 _____ Ulcerative Colitis, Unspecified |

Other: _____

REQUIRED: Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.

LAB RESULTS: Include Negative TB within 12 months.

PRESCRIPTION

Pre-Medications

- Acetaminophen: 650 mg PO
- Cetirizine: 10 mg PO
- Diphenhydramine: 25 mg PO
- Diphenhydramine: 25 mg IVP
- Famotidine: 20 mg PO
- Methylprednisolone: 125 mg SIVP

Other: _____

Lab Orders+

Required: Negative TB, annually

+Medix Infusion will draw maintenance labs unless otherwise directed by Referring Provider

ENTYVIO (vedolizumab)

Loading Dose

IV: Infuse 300 mg in 250 mL of 0.9% Sodium Chloride over at least 30 minutes at weeks 0, 2, 6

Maintenance Dose (SELECT ONE)

- IV:** Infuse 300 mg in 250 mL of 0.9% Sodium Chloride over at least 30 minutes every 8 weeks for one year
- IV:** Infuse 300 mg in 250 mL of 0.9% Sodium Chloride over at least 30 minutes every _____ weeks for one year

Following each infusion, flush with 30 mL 0.9% Sodium Chloride

Is the patient on any other disease modifying therapy? Yes No

If yes, please note therapy and last dose: _____

Post Treatment Observations: The patient is observed for 30 minutes following the first administration.

Adverse Events: In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.

Comments:

PRESCRIBER INFORMATION

Prescriber Name: _____ Signature: _____

Date: _____ NPI #: _____ Specialty: _____

Supervising Physician: _____ (If Applicable)

Address: _____ City: _____ State: _____ Zip: _____

Contact Name: _____ Phone: _____ Fax: _____ Email: _____