

**PATIENT INFORMATION**

Patient Name: \_\_\_\_\_ DOB: \_\_\_\_\_ Phone: \_\_\_\_\_ Sex: M F Ht: \_\_\_\_\_ Wt: \_\_\_\_\_ lbs kg  
Primary Language: \_\_\_\_\_ Allergies: \_\_\_\_\_  
**Patient Preferred Location:** \_\_\_\_\_

&lt;ICD 10 CODE REQUIRED&gt;

**DIAGNOSIS & CLINICAL INFORMATION****ICD 10 Code (PROVIDE COMPLETE CODE)**

ICD 10 Code: \_\_\_\_\_

Description: \_\_\_\_\_

**REQUIRED:** Demographics & Most Recent: H&P, clinical notes, & medication list. Supporting clinical notes to include any past tried and/or failed therapies, intolerance, outcomes, or contraindications to conventional therapy.

**LAB RESULTS:** Include culture report.

**PRESCRIPTION****ORBACTIV (oritavancin)**

Start and follow each infusion with a 10 mL D5W flush

**DO NOT USE Normal Saline for dilution or flushing of IV line as it is incompatible with Orbactiv****Loading Dose**

IV: Infuse 1200 mg in D5W for a total volume of 1000 mL as a single dose over 3 hours

Is the patient on any other disease modifying therapy?    **Yes**    **No**  
If yes, please note therapy and last dose: \_\_\_\_\_

**Post Treatment Observations:** The patient is observed for 30 minutes following the administration.**Adverse Events:** In the event of an adverse reaction occurring at a Medix Infusion suite, utilize the Medix Infusion adverse reactions protocol.**Comments:**\_\_\_\_\_  
\_\_\_\_\_**PRESCRIBER INFORMATION**

Prescriber Name: \_\_\_\_\_ Signature: \_\_\_\_\_

Date: \_\_\_\_\_ NPI #: \_\_\_\_\_ Specialty: \_\_\_\_\_

Supervising Physician: \_\_\_\_\_ (If Applicable)

Address: \_\_\_\_\_ City: \_\_\_\_\_ State: \_\_\_\_\_ Zip: \_\_\_\_\_

Contact Name: \_\_\_\_\_ Phone: \_\_\_\_\_ Fax: \_\_\_\_\_ Email: \_\_\_\_\_