Rituximab Form



Infusion	(rituximab)	FAX TO: 972.499.9210
	PATIENT INFORMATION	
Primary Language:	DOB: Phone: S	·
<icd 10="" code="" required=""></icd>	DIAGNOSIS & CLINICAL INFORMATION	
M31.7 Microscopic Polyangitis	ngitis (Wegener's Granuformatosis GPA)	
	Most Recent: H&P, clinical notes, & medication list. S rapies, intolerance, outcomes, or contraindications t ve Hepatitis B within 3 years. PRESCRIPTION	
Pre-Medications		
Acetaminophen: 650 mg PO Methylprednisolone: 125 mg SIVF	Diphenhydramine: 25 mg IVP Other:	
<u>RITUXIMAB (rituximab)</u>		MEDIX USE ONLY Product to be Used:
	as dictated by patient's insurance* ppropriate product based upon benefit investigation OR (DO NOT SUBSTITUTE) lium Chloride	Rituxan Truxima Ruxience
Loading Dose (SELECT ONE) IV: Infuse 1000 mg	IV: infuse 375 mg/m ² – Required \rightarrow Height:	Weight: Ibs or ka
Frequency and Duration (SELECT Infuse Single Dose Infuse every week for 4 weeks tot Infuse initial dose at day 1 followe Other frequency: Is the patient on any other diseas	TONE) Tal ad by 2nd dose on day 15, then repeat cycle every for one year e modifying therapy? Yes No	
	st dose: patient is observed for 60 minutes following the first administ	ration
	f an adverse reaction occurring at a Medix Infusion suite, utiliz	
	PRESCRIBER INFORMATION	
Prescriber Name:	Signature:	
	Specialty:	
Supervising Physician:		(If Applicable)
Address:	City:	State: Zip:

Contact Name: ______ Phone: _____ Fax: _____ Email: _____

medix